

Effect of Cognitive Ability on Oral Health Status: A Scoping Review of Observational Studies

Bidisha Ray^{1*} Saudamini More² Parth Ghatalia³ Tejal Patil⁴ Anamika Shinde⁵ Suyog Savant⁶

¹BDS, Department of Public Health Dentistry, Bharati Vidyapeeth Dental College & Hospital, Navi Mumbai, India.

²MDS, Department of Public Health Dentistry, Bharati Vidyapeeth Dental College & Hospital, Navi Mumbai, India.

³BDS, Department of Public Health Dentistry, Bharati Vidyapeeth Dental College & Hospital, Navi Mumbai, India.

⁴MDS, Department of Oral & Maxillofacial Surgery, Bharati Vidyapeeth Dental College & Hospital, Navi Mumbai, India.

⁵BDS, Department of Oral & Maxillofacial Surgery, Bharati Vidyapeeth Dental College & Hospital, Navi Mumbai, India.

⁶MDS, Department of Public Health Dentistry, Bharati Vidyapeeth Dental College & Hospital, Navi Mumbai, India.

ABSTRACT

Background: This review article aims to update and consolidate the scientific data available regarding the link between cognition and oral health in the elderly population. This review was undertaken in May 2020. A rigorous data search and selection strategy in accordance with the PRISMA guidelines was employed. The population (P) considered was 'elderly' (more than 50 years of age), exposure (E) was cognitive difficulty, compared (C) with healthy individuals (in few studies) and the outcome (O) assessed was oral health status. English language articles published from January 1990 to May 2020 were searched. Twenty five articles were identified that assessed oral health and at least one of the specific cognitive domains along with at least one cofactor in an older adult cohort study. For every cognitive domain like learning and memory, complex attention and executive function most studies identified significant associations with oral health (when covariates were not adjusted for). In addition, clinical determined poorer oral health status was associated with less frequent of dental visits. In presence of cardiovascular disease and inflammatory markers in geriatric patients, oral health status has shown to be poor. The present review incorporated longitudinal, prospective cohort studies along with cross sectional studies. This review mediates evidence of an association between cognitive functions with oral health in old age along with other cofactors like associated systemic disease and socioeconomic status. The present article will help clinicians better understand the role of cognitive functions in oral health conditions in elderly.

Keywords: Geriatric Dentistry; Periodontal Health; Edentulous.

INTRODUCTION

Cognition is critical for functional independence as people age, irrespective of whether a person is independent or dependent. Cognitive abilities can be divided into several specific cognitive domains including attention, memory, executive cognitive function, language, and visuospatial abilities. Prospective memory, specifically remembering to perform intended action in the future (e.g., taking medication after breakfast). Whereas, executive cognitive function is particularly important for

novel tasks. These are not associated with a habitual responses necessarily, but depends critically on the prefrontal cortex.^[1] Cognitive dementia, according to WHO, is a syndrome in which there is deterioration in memory, thinking, behaviour, and the ability to perform everyday activity. According to a report published by WHO in September 2019, around 50million of the population have dementia.^[2] This is the major reason behind dependency and disabilities during the old age.

Received: Jan. 3, 2021: Accepted: Mar. 2, 2021

*Correspondence Dr. Saudamini More.

Department of Public Health Dentistry, Bharati Vidyapeeth Dental College & Hospital, Navi Mumbai, India.

Email: drsaudamini90@gmail.com

As per the last census report (2011), 104 million (53million female & 51 million male) of population belong to the senior citizens in India. Ageing is a natural biological process which is primarily influenced by our genetic makeup. However, in recent centuries it has become more heavily influenced by social, environmental, economic and medical changes as well as physical and mental health [3]. Age related disabilities and immunodeficiency is related to under-nutritional status in the population below the poverty line.[4]

Sir Bernard Isaacs coined the term 'Geriatric Giant' in the 1960s to describe categories of impairments in the elderly. Falls were associated with frailty which in turn were associated with neurocognitive decline and delirium as an older person became less agile and in deteriorating health.[3] As dental and oral health issues are a newly recognized burden on the health of the elderly, only scant regard for the implications of overt dental caries and active periodontal infection has occurred.[5] However, as health issues are interconnected, maintaining poor oral health can have consequences that are not immediately evident.

Thus, the present review aims to understand & update the scientific literature regarding the link between cognition and oral health in the elderly population. Different cofactors like cardiovascular disease , diabetes, lifestyle habits are also considered and their roles in the deterioration of the oral health is mentioned.

METHODOLOGY

The PRISMA guidelines were followed for a methodological approach to include and various articles from the relevant databases. Various articles based on topics of cognitive dementia, oral health and aging were assessed and screened.

A systematic literature search was carried out in the following databases: Pubmed , WHO ,Medline and ScienceDirect. The keywords used were 'cognitive psychology', 'Alzheimer' , 'diabetes', 'ageing', 'oral health', 'periodontal therapy ,cardiovascular disease, dementia, attitude, tooth loss, denture problem, dry mouth. The filter applied in the search included the language (English) and date of publication (1990–2020).A final search was carried out on 28 May 2020 to ensure the most recent

literature was included in this review. In addition, the reference lists and bibliographies of all relevant studies were looked for additional sources. The results from the search were organised and duplicate references were removed using the Mendeley bibliographic software. The title and abstract of the remaining studies were assessed by authors independently using the inclusion and exclusion criteria for suitability. In the case where it was difficult to make a decision on the basis of the title and abstract only, the full text was obtained for further assessment. A total of 25 studies which met the inclusion criteria and were included in the review. Figure 1 depicts PRISMA flowchart for the selection strategy employed for selecting studies.

Inclusion and exclusion criteria

- 1) The selected databases are based on Cohort studies that have studied the cognitive function of the older age group (>50years) by standardized DSST (Digital Symbol Substitution test), Mini-Mental State Examination (MMSE) , **Delayed word recall (DWR) and Word fluency (WF).**
- 2) Covariates have been considered like associated disease like Cardiovascular disease and diabetes, mental health issue like depression , socioeconomic status and education background, nutrition, demography. The participants were either dependant or community dwellers.
- 3) Oral health assessment scales like OHI-S Index , Periodontal Index, DMFT Index, self-assessment and dental follow-ups at specific time span were included
- 4) The date of publication of the articles were between 1990-2020. The study designs were Cohort cross-sectional and/or longitudinal with at least one above mentioned covariates.

Data sources ,search strategy and study selection

Data extraction

A data extraction form was developed and piloted independently by two authors and modified as required . The information including author, year of publication, country, characteristics of

sample/questionnaire and key outcome items was extracted by two authors and checked by one author for accuracy and completeness of the results. There was clear heterogeneity among the studies in their approaches to measuring and reporting the

knowledge, attitudes, and practices in reference to oral health care. Therefore descriptive analysis was administered in most sections and data was pooled to calculate mean percentage wherever studies had similar outcome items to match.

Fig 1: PRISMA Flow diagram depicting selected studies




Table 1: Data extracted from various articles.

Article	Sample Size (n)	Study Design	Co-variates	Result
---------	-----------------	--------------	-------------	--------

Kweider M et al. (1993) [6]	(n=50)	Cross-sectional	Associated Cardiac Disease Age (>50)	Fibrinogen and WBC count was established as a mediator between cardiac and inflammatory dental disease.
Saxe et al. (1995) [7]	(n=129)	Cross-section	Demography Age (>50)	Cavity filling using amalgam has no impact on cognitive impairment.
Cohen-Mansfield J et al (2004) [8]	(n=80)	Cross-sectional	Age (>59)	The findings reveal that the MDS (Minimum Data Set, a comprehensive functional assessment) underreports pain in cognitively impaired residents. Participants reported excruciating pain of oral soft tissue lesions and decreased pain threshold.
Kim JM et al(2006) [4]	(n= 1,204)	Cross-sectional	Nutritional status ;chronic inflammatory condition Age (>65)	Fewer teeth were associated with and increased risk of incident dementia along with marked reduction in nutritional status.
Wu B et al. (2007) [9]	(n= 1984)	Cross-sectional	Socio-demography Associated disease Age (>60)	Higher cognitive level & higher sociodemographic level were associated with frequent dental visit Poor health status is associated with poor oral health
Gagliardi D et al 2008 [10]	(n= 253)	Longitudinal	Socioeconomic condition Age (> 75)	Oral health related quality of life improved with increased state funded health care.
Yu and Kuo (2008) [11]	(n=803)	Cross-sectional	Demography Associated disease Age (>60)	Better periodontal health is associated with higher DSST scores (unadjusted).
Lexomboon D et al (2012) [12]	(n= 557)	Cross-sectional	Associated disease Age (>77)	Adjusted for sex, age, education, depression, and mental illness, odds of cognitive impairment significantly higher for persons with chewing difficulty
Naorungroj et al. (2013) [13]	(n= 10,050)	Longitudinal	Age, socioeconomic condition, demography, habit, associated disease. Age (> 56)	Tooth loss and periodontal problems shows close association with decline in memory domain of cognition.
Stewart et al. (2013) [14]	(n= 1,053)	Cross-sectional	Age, Demography, Poor mental health. Age (>70)	Decline complex attention is associated with poor oral health. Lower DSST scores are associated with poor Gingival Index results.
Asai K et al. (2015) [15]	(n= 8,124)	Cross-sectional	Associated disease (Cardiac Disease, diabetes, etc)	The study suggests a linear relationship between tooth loss and degree of arterial stiffness that is sex dependent.

Foltyn P et al (2015) [3]	(n=8)	Longitudinal	Age (>75) Age, habit, socioeconomic condition Age (>50)	The contributing factors for poor oral health & reduced motor function, declining cognition lead to an increase in both morbidity and mortality also impacting the quality of life.
Tsakos G et al (2015) [16]	(n=3,166)	Longitudinal	Depression, socioeconomic condition and Comorbidities Age (>60)	Tooth loss is a potential early marker of decline in older age. Edentulous participants recalled 0.88% less words at 0.09 m/s slower speed than dentulous participants (adjusted for time & demography)
Supawadee Naorungroj et al 2015 [17]	(n=911)	Prospective cohort study	Cardiovascular Disease Age (>52)	Mean decline in scores of Digit Symbol Substitution Test (DSST), Delayed word recall (DWR) and Word fluency (WF) was associated with tooth loss and poor Periodontal health.
Kamer AR et al (2016) [18]	(n=152)	Cross-sectional	Alzheimer's Disease Periodontal inflammation Age (> 60)	Participants with existing periodontal inflammation showed lower Digital Symbol test scores. Also no. of missing teeth were more.
Komiyama T et al (2016) [19]	(n=834)	Prospective cohort study	Habit, depression, Education Age (>70)	Participants with less than 20 teeth showed higher chances of developing functional disability. Thus, tooth loss was associated is greater risk of functional disability.

Table 2: Data extracted from various articles

Article	Sample Size (n)	Study design	Co-variates	Results
Kundapur V et al (2017) [20]	(n=141)	Cross-sectional	Demography Age (>60)	With loss of multiple teeth and declining motor skills prosthetic denture wear is difficult to adapt. Leading to rapid bone loss.
Juan Li et al. 2017 [21]	(n= 8,153)	Longitudinal	Socioeconomic Condition Age (> 60)	More teeth were associated with better cognitive function. Mean baseline teeth number-17.5, the mean of baseline cognitive function-27.3
Takada et al. (2018) [22]	(n=64)	Oral Health Intervention	Age (>65)	The cognitive test scores showed positive results after regular oral health intervention.
Saito S et al. (2018) [23]	(n=140)	Longitudinal	Associated disease (Cardiac Disease, diabetes) Age (>65)	Multiple tooth loss significantly increased the risk of developing cognitive impairment. Association between Chronic inflammatory systemic disease and periodontal disease. Mini-Mental State Examination (MMSE) score were less in participants with 0-9 teeth in oral cavity.
Yoo JJ et al. (2019) [24]	(n= 2,09,806)	Longitudinal	Dementia, Demography	Individuals with tooth loss had a higher risk for dementia than those without

			Age (> 60)	tooth loss. Incidence of dementia decreased with periodontal treatment.
Kim DW et al. (2019) [25]	(n=10,215)	Retrospective Cohort study	Diabetes Age (>65)	Diabetic individuals had a higher risk of tooth loss, due to periodontal problem than nondiabetic individuals (HR = 1.298)
• Kato H et al. (2019) [26]	(n= 210)	Longitudinal	Objective Dementia Age (>60)	Individuals with a total of 20 teeth or more, including both natural and artificial teeth, had significantly higher MMSE scores
Thomson et al (2019) [27]	(n=818)	Prospective cohort study	Socioeconomic Condition, education Age (> 50)	IQ test scores were highest with better DMFT Index and OHI-S Index.
Gao SS et al (2020)[28]	(n=341)	Cross-sectional	Age (>65) Demography	Around 129 participants with dementia showed poor DMFT score and poor toothbrushing habits.

RESULT

Table 1 & 2 show the results of various studies which were reviewed. Social cognitive theory and oral health implication have presented its results in 1 study, with covariates of socioeconomic background and tobacco smoking. Covariates like associated disease and risk factor, education and socioeconomic background, smoking and tobacco abuse, nutrition have been considered. Various articles published from the year 1990 to 2020 with a sample size ranging from a minimum of n= 8 to maximum of n=2,09,806 and participant age ranged 50years and above were compiled.

Cognitive tests using Digit Symbol Substitution Test (DSST) (12 articles) and standardized mini-mental state examination (SMMSE) (10 articles) were used. Cognitive functions of key 6 domains, according to the DSM-5 classification system, were considered for the examination. However, all the 25 articles established a link between cognitive domains of learning & memory (16 articles), executive function (13 articles), complex attention (8 articles) and their implication on the deterioration of oral health.

The participants selected in 23 studies had their oral health assessed by examinations that were conducted in a time interval of 6 months to a year in each of the studies. Periodontal health assessment and number of missing teeth was recorded and considered as standard.

Self- assessment technique was used for 1 study by preparing a questionnaire according to the GOHAI (Geriatric Oral Health Assessment Index) survey. Sample size (n=141) were subjected to 16 questions (4 on physical functions, 5 on psychological, 3 on pain and discomfort) with consent.

DISCUSSION

Chronic periodontitis is a widespread peripheral immune-inflammatory condition. It is postulated to play an important role in the aggravation of Dementia. An association between peripheral inflammation and dementia was hypothesized by an article published by Jae-Min Kim et al in the International Journal of Geriatric Psychiatry (2007). It was suggested that chronic systemic inflammation aggravated the inflammatory response in the brain. This can be due to increase in the levels of pro-inflammatory mediators such as tumour necrosis factor- α , interleukin-1, interleukin -6 and in the plasma.^[5] The contributing factors for poor oral health like dental caries, acute and chronic periodontal infections and compromised systemic health causing xerostomia, reduced fine motor function^[6]. Thus, the declining cognition and motivation may not only lead to both morbidity and mortality but also impact on quality of life.

Associated health problems like cardiovascular disease, diabetes and other chronic inflammatory systemic disease have shown unfavourable outcomes like plaque accumulation, periodontal

disease followed by tooth loss. Oral health in later life is one of the significant determinants of quality of life (QoL) and life satisfaction as well as a key health status indicator related to functional limitations and limited daily activities.^[27] When elderly become care dependent, their oral health often deteriorates due to neglect of oral (self)care. Socioeconomic factor also plays a key factor amongst this population that forms a contributing factor. The lack of attention for oral care can lead to poor oral health and can be considered a hidden health hazard as dental awareness and oral health has been shown to significantly contribute to general health and quality of life.^[29,30]

With age, changes in biological and physiological aspects are inevitable. There are ample of evidences that shows alterations in brain structure and performance are intimately tied to alterations in cognitive function. Crystallized intelligence refers to skills, ability, and knowledge that's overlearned, well-practiced, and familiar. Crystallized abilities like vocabulary and general knowledge remain stable or gradually improve at a rate of 0.02 to 0.003 standard deviations per year through the sixth and seventh decades of life. . Fluid cognition is a person's natural ability to process and learn new knowledge, solve problems, attend to and control one's environment. Post the third decade of life i.e the peak, fluid cognition, especially psychomotor ability and processing speed declines at an estimated rate of -0.02 standard deviations per year.^[31]

Just as age-related changes in brain structure and performance aren't uniform across the entire brain or across individuals, age-related changes in cognition aren't uniform across all cognitive domains or across all older individuals. The basic cognitive functions most suffering from age are attention and memory. Attention is that the means by which we actively process a limited amount of data from the big amount of data available through our senses, our stored memories, and our other cognitive processes. A decline in the domain of attention can have widespread effects on one's ability to function adequately and efficiently in their daily life.

The prevalence of cardiovascular disease (CVD) is increasing rapidly in the growing aging population. Cardiovascular disease contributes to

morphological alteration of the brain that leads to cognitive decline, dementia and cerebrovascular disease. The cognitive decline occurs because of the formation of amyloid deposits that provoke neuro-inflammation, finally resulting in necrobiosis^[32]. Associations between high blood pressure (BP) and poor performance on neuropsychological tests are well described in the neuropsychological literature particularly in the areas of memory, attention, and executive function. Metabolic-related conditions such as diabetes and obesity have also been associated with cognitive impairment. Periodontal disease, a local inflammatory process is a common oral problem of the older population leading to loss of natural teeth. This local inflammatory process may induce a systemic inflammatory reaction. Therefore, periodontitis could trigger and/or exacerbate an inflammatory condition especially in elderly subjects, resulting in memory impairments, contributing to accelerate the progression of neurodegenerative diseases.

Each of these articles, through cross section analysis (10 studies) and longitudinal studies (8 studies) have established satisfactory results supporting the hypothesis positively. However, none of the articles has considered all the covariates altogether. The database was collected through National health surveys or self-reported. Thus, the uniformity in the measurement is lacking.

CONCLUSION

This scoping review mediates evidence of an association between cognitive functions with oral health in old age along with other cofactors like associated systemic disease and socioeconomic status. Thus, understanding that life expectancy is impacted by systemic health and the 'Geriatric Giants', the dental professional's role is to make the individual comfortable and maintain function as best possible. Elderly with complex care needs experience significantly more oral health problems and are more often edentulous than robust elderly. Moreover, elderly with few teeth and elderly with implant-supported overdentures reported less frailty and better quality of life. .However, frequent oral health check-ups for dependent as well as independent aged population can be incorporated for better oral health status. Policies at community level for population of lower socioeconomic background can achieve better quality of life.

CONFLICTS OF INTEREST

The authors declare they have no potential conflict of interest regarding this article.

REFERENCES

1. Sternberg RJ, Sternberg K. Memory: Models and research methods. *Cognitive Psychology*. 2012;185-227
2. World Health Organization. [www. who. int/news-room/fact-sheets/detail/dementia](http://www.who.int/news-room/fact-sheets/detail/dementia).
3. Foltyn P. Ageing, dementia and oral health. *Australian dental journal*. 2015 Mar;60:86-94.
4. Kim JM, Stewart R, Prince M, Kim SW, Yang SJ, Shin IS, Yoon JS. Dental health, nutritional status and recent-onset dementia in a Korean community population. *International Journal of Geriatric Psychiatry: A journal of the psychiatry of late life and allied sciences*. 2007 Sep;22(9):850-5.
5. Gaur S, Agnihotri R. A lzheimer's disease and chronic periodontitis: Is there an association?. *Geriatrics & gerontology international*. 2015 Apr;15(4):391-404.
6. Kweider M, Lowe GD, Murray GD, Kinane DF, McGowan DA. Dental disease, fibrinogen and white cell count; links with myocardial infarction?. *Scottish medical journal*. 1993 Jun;38(3):73-4.
7. Saxe SR, Snowdon DA, Wekstein MW, Henry RG, Grant FT, Donegan SJ, Wekstein DR. Dental amalgam and cognitive function in older women: findings from the Nun Study. *The Journal of the American Dental Association*. 1995 Nov 1;126(11):1495-501.
8. Cohen-Mansfield J. The adequacy of the minimum data set assessment of pain in cognitively impaired nursing home residents. *Journal of Pain and Symptom Management*. 2004 Apr 1;27(4):343-51.
9. Wu B, Plassman BL, Liang J, Wei L. Cognitive function and dental care utilization among community-dwelling older adults. *American Journal of Public Health*. 2007 Dec;97(12):2216-21.
10. Gagliardi D, Slade GD, Sanders AE. Impact of dental care on patients' treatment goals and quality of life. *Aust Dent J*. 2008;53(1):26-33.
11. Yu YH, Kuo HK. Association between cognitive function and periodontal disease in older adults. *Journal of the American Geriatrics Society*. 2008 Sep;56(9):1693-7.
12. Lexomboon D, Trulsson M, Wårdh I, Parker MG. Chewing ability and tooth loss: association with cognitive impairment in an elderly population study. *Journal of the American Geriatrics Society*. 2012 Oct;60(10):1951-6.
13. Naorungroj S, Slade GD, Beck JD, Mosley TH, Gottesman RF, Alonso A, Heiss G. Cognitive decline and oral health in middle-aged adults in the ARIC study. *Journal of dental research*. 2013 Sep;92(9):795-801.
14. Stewart R, Weyant RJ, Garcia ME, Harris T, Launer LJ, Satterfield S, Simonsick EM, Yaffe K, Newman AB. Adverse oral health and cognitive decline: the health, aging and body composition study. *Journal of the American Geriatrics Society*. 2013 Feb;61(2):177-84.
15. Asai K, Yamori M, Yamazaki T, Yamaguchi A, Takahashi K, Sekine A, Kosugi S, Matsuda F, Nakayama T, Bessho K, Nagahama Study Group. Tooth loss and atherosclerosis: the Nagahama Study. *Journal of Dental Research*. 2015 Mar;94(3_suppl):52S-8S.
16. Tsakos G, Watt RG, Rouxel PL, de Oliveira C, Demakakos P. Tooth loss associated with physical and cognitive decline in older adults. *Journal of the American Geriatrics Society*. 2015 Jan;63(1):91-9.
17. Naorungroj S, Schoenbach VJ, Wruck L, Mosley TH, Gottesman RF, Alonso A, Heiss G, Beck J, Slade GD. Tooth loss, periodontal disease, and cognitive decline in the Atherosclerosis Risk in Communities (ARIC) study. *Community dentistry and oral epidemiology*. 2015 Feb;43(1):47-57.
18. Kamer AR, Fortea JO, Videla S, Mayoral A, Janal M, Carmona-Iragui M, Benejam B, Craig RG, Saxena D, Corby P, Glodzik L. Periodontal disease's contribution to Alzheimer's disease

- progression in Down syndrome. *Alzheimer's & Dementia: Diagnosis, Assessment & Disease Monitoring*. 2016 Jan 1;2:49-57.
19. Komiyama T, Ohi T, Miyoshi Y, Murakami T, Tsuboi A, Tomata Y, Tsuji I, Watanabe M, Hattori Y. Association between tooth loss, receipt of dental care, and functional disability in an elderly Japanese population: the Tsurugaya project. *Journal of the American Geriatrics Society*. 2016 Dec;64(12):2495-502.
20. Kundapur V, Hegde R, Shetty M, Mankar S, Hilal M. Effect of loss of teeth and its association with general quality of life using geriatric Oral health assessment index (Gohai) among older individuals residing in rural areas. *International journal of biomedical science: IJBS*. 2017 Mar;13(1):6.
21. Li J, Xu H, Pan W, Wu B. Association between tooth loss and cognitive decline: a 13-year longitudinal study of Chinese older adults. *PloS one*. 2017 Feb 3;12(2):e0171404.
22. Tominaga T, Tomotake M, Takeda T, Ueoka Y, Tanaka T, Watanabe SY, Kameoka N, Nakataki M, Numata S, Izaki Y, Sumitani S. Relationship between social and cognitive functions in people with schizophrenia. *Neuropsychiatric disease and treatment*. 2018;14:2215.
23. Saito S, Ohi T, Murakami T, Komiyama T, Miyoshi Y, Endo K, Satoh M, Asayama K, Inoue R, Kikuya M, Metoki H. Association between tooth loss and cognitive impairment in community-dwelling older Japanese adults: a 4-year prospective cohort study from the Ohasama study. *BMC Oral Health*. 2018 Dec;18(1):1-8.
24. Yoo JJ, Yoon JH, Kang MJ, Kim M, Oh N. The effect of missing teeth on dementia in older people: a nationwide population-based cohort study in South Korea. *BMC oral health*. 2019 Dec 1;19(1):61.
25. Yoo JJ, Kim DW, Kim MY, Kim YT, Yoon JH. The effect of diabetes on tooth loss caused by periodontal disease: A nationwide population-based cohort study in South Korea. *Journal of Periodontology*. 2019 Jun;90(6):576-83.
26. Kato H, Takahashi Y, Iseki C, Igari R, Sato H, Sato H, Koyama S, Tobita M, Kawanami T, Iino M, Ishizawa K. Tooth loss-associated cognitive impairment in the elderly: a community-based study in Japan. *Internal Medicine*. 2019:1896-.
27. Thomson WM, Broadbent JM, Caspi A, Poulton R, Moffitt TE. Childhood IQ predicts age-38 oral disease experience and service-use. *Community dentistry and oral epidemiology*. 2019 Jun;47(3):252-8.
28. Ren T, Gao Y, Qiu Y, Jiang S, Zhang Q, Zhang J, Wang L, Zhang Y, Wang L, Nie K. Gut microbiota altered in mild cognitive impairment compared with normal cognition in sporadic Parkinson's disease. *Frontiers in neurology*. 2020 Feb 25;11:137.
29. Li X, Kolltveit KM, Tronstad L, Olsen I. Systemic diseases caused by oral infection. *Clinical microbiology reviews*. 2000 Oct 1;13(4):547-58.
30. Baiju RM, Peter EL, Varghese NO, Sivaram R. Oral health and quality of life: current concepts. *Journal of clinical and diagnostic research: JCDR*. 2017 Jun;11(6):ZE21.
31. Harada CN, Love N, MC T. KL (2013). Normal cognitive aging. *Clinics in Geriatric Medicine*.;29(4):737-52.
32. Leritz EC, McGlinchey RE, Kellison I, Rudolph JL, Milberg WP. Cardiovascular disease risk factors and cognition in the elderly. *Current cardiovascular risk reports*. 2011 Oct 1;5(5):407.