

Methods of Canine Retraction in Orthodontics: Traditional to Conventional Approaches

Apurva Chavan^{1*}, Amit Handa¹, Himanshu Srivastava², Dipali Mane¹, Aditi Sarda³

¹Department of Orthodontics and Dentofacial Orthopaedics, Aditya Dental College and Hospital, Beed, Maharashtra, India.

²Department of Oral and Maxillofacial Pathology and Microbiology, Aditya Dental College and Hospital, Beed, Maharashtra, India.

³Department of Conservative Dentistry and Endodontics, Aditya Dental College and Hospital, Beed, Maharashtra, India.

Abstract

Background: Orthodontic treatment frequently necessitates the retraction of canines to correct dental malocclusions and achieve optimal occlusion. This review examines a range of methods for canine retraction in orthodontics, covering both traditional techniques and contemporary approaches. Traditional methods, such as elastic chains, NiTi coil springs, and power arms, provide effective and economical solutions but may require frequent adjustments and can cause patient discomfort. Contemporary methods, including temporary anchorage devices, clear aligner therapy, self-ligating brackets, and piezoelectric surgery, offer enhanced control, efficiency, and patient comfort. These newer techniques often involve advanced materials and technology, providing more predictable and efficient tooth movements. Each method's mechanism, advantages, and limitations are discussed in detail to offer a comprehensive understanding of their clinical applications. This review highlights the importance of selecting the appropriate method based on individual patient needs, the severity of malocclusion, and clinical expertise. The evolution of canine retraction methods underscores the advancements in orthodontic treatment, aiming to improve patient outcomes and treatment efficiency. Future research and technological developments will continue to shape the landscape of orthodontics, offering even more refined and effective solutions for canine retraction.

Keywords: Canine retraction, Contemporary methods, Dental malocclusion, Orthodontic appliances, Orthodontics, Traditional methods.

INTRODUCTION

Orthodontic treatment aims to align teeth, improve occlusion, and enhance esthetic appearance. One critical aspect of orthodontic therapy is the retraction of canines, particularly in cases involving premolar extraction, to correct dental malocclusions. Malocclusions, which are misalignments of teeth or incorrect relations between the teeth of the two dental arches, affect a significant portion of the population and can lead to various functional and esthetic issues.^[1-3] Effective canine retraction is essential in many orthodontic treatment plans as it helps achieve proper occlusion and enhances facial esthetics.

IMPORTANCE OF CANINE RETRACTION

Canine retraction is often necessary in orthodontic cases where space needs to be created for the alignment of other teeth, particularly when premolars have been extracted. The movement of canines into the extraction sites allows for the correction of crowding and the establishment of an optimal bite. Canines play a crucial role in the dental arch due to their

long roots and strategic position, which contributes to the overall stability and function of the dentition.^[2-5] Achieving efficient and controlled retraction of canines is therefore pivotal in many orthodontic treatments.

EVOLUTION OF CANINE RETRACTION METHODS

Over the years, the methods used for canine retraction have evolved significantly. Traditional techniques, while still in use, have been supplemented by more contemporary approaches that leverage advances in materials and technology. The goal of these advancements is to enhance the predictability, efficiency, and comfort of orthodontic treatments.^[6-10]

Address for correspondence: Dr. Apurva Chavan,
Department of Orthodontics and Dentofacial Orthopaedics,
Aditya Dental College and Hospital, Beed, Maharashtra, India.
Email: arnavarvichavan21@gmail.com

Received: Aug. 9, 2024; **Accepted:** Sept. 29, 2024; **Published:** Oct. 15, 2024

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

How to cite this article: Chavan A, Handa A, Srivastava H, Mane D, Sarda A. Methods of Canine Retraction in Orthodontics: Traditional to Conventional Approaches. J Res Adv Dent 2025;16(1):1-5.

Access this article online

Website:
www.jrad.co.in

DOI:
<https://doi.org/10.53064/jrad.2024.16.1.636>

TRADITIONAL METHODS OF CANINE RETRACTION

Elastic chains and elastics

Elastic chains and elastics have long been a staple in orthodontic treatment, particularly for the retraction of canines. These materials are favored for their simplicity, cost-effectiveness, and ease of use. They are an integral part of traditional orthodontic techniques aimed at correcting dental malocclusions and achieving optimal alignment of the dental arches.^[1-5,11]

Mechanism

Elastic chains and elastics function by applying a continuous pulling force on the teeth, specifically targeting the canines for retraction. Elastic chains are made of a series of elastic loops linked together, forming a chain that can be stretched and attached to brackets on the teeth. Elastics, on the other hand, are single rubber bands that can be stretched between brackets or hooks on the braces. When these elastics or chains are stretched, they exert a pulling force on the teeth, gradually moving them into the desired position.

The principle behind their use is based on the application of orthodontic forces that stimulate bone remodeling. The tension created by the elastics or chains causes pressure on the periodontal ligament and the alveolar bone, leading to resorption on the pressure side and deposition on the tension side. This biological response allows for the gradual movement of the teeth through the bone.

Clinical application

Elastic chains and elastics are typically used in conjunction with fixed orthodontic appliances, such as braces. They are attached to brackets or hooks on the braces to create the desired retraction force. The orthodontist will determine the appropriate length and tension based on the specific treatment plan and the patient's needs.

For canine retraction, elastic chains may be attached from the canine brackets to the posterior teeth or to auxiliary hooks placed on the archwire. This setup allows for the controlled retraction of the canines into the spaces created by premolar extractions. The tension in the elastic chains must be carefully monitored and adjusted to ensure consistent force application and prevent excessive pressure that could damage the teeth or supporting structures.

In summary, elastic chains and elastics are valuable tools in traditional orthodontic treatment for canine retraction. Their simplicity, cost-effectiveness, and ability to provide continuous force make them a popular choice among orthodontists. However, their limitations, such as loss of elasticity, frequent replacements, and potential for discomfort, must be carefully managed to ensure effective and efficient treatment outcomes. By understanding the mechanisms, advantages, and limitations of elastic chains and elastics, orthodontists can make informed decisions about their use in clinical practice.

Nickel-titanium (NiTi) coil springs

NiTi coil springs have become a fundamental component in orthodontic treatments due to their unique properties of superelasticity and shape memory. These springs are particularly useful for canine retraction, providing a continuous and consistent force over an extended period, which is essential for effective tooth movement.^[12-15]

Mechanism

NiTi coil springs are made from a NiTi alloy known for its superelastic and shape memory characteristics. These properties allow the springs to return to their original shape after being deformed. In orthodontics, NiTi coil springs are attached to brackets and exert a continuous force on the teeth to be moved. The force generated by these springs remains relatively constant, regardless of the amount of activation or deformation, making them highly effective for sustained tooth movement.

When NiTi coil springs are activated, they apply a gentle yet consistent force to the teeth, stimulating the periodontal ligament and alveolar bone to remodel. This biological process involves bone resorption on the pressure side and bone deposition on the tension side, allowing the teeth to move gradually into their desired positions.

Clinical application

NiTi coil springs are typically used in conjunction with fixed orthodontic appliances, such as braces. They can be attached between the brackets of the teeth to be retracted and the posterior teeth or anchorage points. The orthodontist activates the springs by stretching them to the desired length, which then exerts a continuous retraction force on the canines.

For canine retraction, NiTi coil springs are often preferred over elastic chains due to their ability to maintain consistent force over a longer period. This consistent force is crucial for efficient and controlled movement of the canines into the extraction spaces, ensuring proper alignment and occlusion.

In clinical practice, NiTi coil springs can be used in various configurations depending on the treatment goals and the patient's specific needs. For example, they can be placed directly on the archwire or attached to auxiliary hooks or power arms to achieve the desired force vectors and tooth movements. The orthodontist must regularly monitor the springs to ensure they are functioning correctly and making the necessary adjustments to optimize treatment outcomes.

Power arms and auxiliary wires

Power arms and auxiliary wires are integral components in orthodontic treatment, particularly in cases requiring precise and controlled movement of the canines. These devices enhance the efficiency of tooth movement by optimizing the application of orthodontic forces, thereby reducing tipping and promoting bodily movement. Their strategic use in conjunction with fixed appliances facilitates better control over the direction and magnitude of forces applied to the teeth.^[16-20]

Mechanism

Power arms are small extensions attached to the orthodontic brackets, typically on the canines or premolars. They serve as leverage points, allowing for the application of forces closer to the tooth's center of resistance. This proximity reduces tipping tendencies and promotes more uniform bodily movement. Auxiliary wires, often referred to as piggyback wires or bypass wires, are additional archwires used to guide and control tooth movement. These wires are threaded through the brackets and connected to the power arms, ensuring that the forces are directed as needed for effective retraction.

The combined use of power arms and auxiliary wires allows for a multi-vector approach to force application. This means that forces can be adjusted in various directions and magnitudes, providing a higher level of control over tooth movement. The precise placement and activation of these components are crucial for achieving the desired orthodontic outcomes.

Clinical application

Power arms and auxiliary wires are typically used in conjunction with fixed orthodontic appliances. The orthodontist attaches power arms to the brackets on the canines or adjacent teeth. Auxiliary wires are then threaded through the brackets and connected to the power arms, creating a system that allows for precise force application.

For canine retraction, power arms are often positioned to extend the point of force application, bringing it closer to the center of resistance of the canines. This setup reduces the tipping tendency and promotes bodily movement. Auxiliary wires can be used to distribute forces evenly across the dental arch, ensuring that the retraction forces are directed appropriately.

CONTEMPORARY METHODS OF CANINE RETRACTION

Temporary anchorage devices (TADs)

TADs have emerged as a revolutionary tool in orthodontics, offering unparalleled control and precision in tooth movement. These devices, also known as mini-implants or microimplants, provide absolute anchorage, enabling orthodontists to achieve complex movements that were previously difficult or impossible with traditional methods. TADs are particularly beneficial for canine retraction, providing a stable anchorage point that minimizes undesired tooth movements and enhances treatment predictability.^[19-22]

Mechanism

TADs are small, screw-like devices made from biocompatible materials, such as titanium alloys. They are temporarily inserted into the alveolar bone, usually under local anesthesia, to serve as stable anchorage points for applying orthodontic forces. Once in place, TADs can be used to apply forces directly to the teeth, eliminating the need for traditional anchorage methods, such as headgear or intraoral appliances, which often rely on patient compliance and cooperation.

The stability of TADs allows for the application of forces in multiple directions, providing a high degree of control over tooth movement. This is particularly useful for canine retraction, where precise control is necessary to achieve the desired alignment without affecting adjacent teeth.

Clinical application

TADs are typically placed in the alveolar bone in strategic locations to provide optimal anchorage for canine retraction. The orthodontist determines the most appropriate placement sites based on the individual patient's dental and skeletal anatomy. Common sites for TAD placement include the interradicular spaces between teeth, the palate, and the retromolar area.

Once placed, TADs can be immediately loaded with orthodontic forces. For canine retraction, TADs are often used in combination with elastomeric chains, coil springs, or other orthodontic appliances to apply the desired retraction force. The precise control offered by TADs allows for efficient and predictable movement of the canines into the extraction spaces.

Clear aligner therapy

Clear aligner therapy has gained significant popularity in recent years as a modern, esthetic alternative to traditional braces. This method involves the use of custom-made, transparent plastic trays that fit snugly over the teeth and gradually move them into their desired positions. Clear aligners are particularly appealing due to their removability, improved esthetics, and increased comfort compared to conventional fixed appliances. This section explores the mechanisms, advantages, limitations, and clinical applications of clear aligner therapy, with a focus on its role in canine retraction.^[16,18,23,24]

Mechanism

Clear aligners are designed using advanced digital technology. The treatment begins with a comprehensive assessment of the patient's dental condition, including digital scans, photographs, and radiographs. These records are used to create a detailed 3D model of the patient's teeth, which serves as the basis for the treatment plan. Specialized software then simulates the sequential movement of teeth, creating a series of aligners, each slightly different from the last, to gradually guide the teeth into the desired positions.

For canine retraction, clear aligners apply gentle, controlled forces to the canines, encouraging them to move into the extraction spaces created by the removal of premolars. The aligners are worn for approximately 20–22 h per day and are replaced with the next set in the series every 1–2 weeks, depending on the treatment plan. This gradual, step-by-step approach ensures steady progress and minimizes discomfort.

Clinical application

Clear aligner therapy is initiated with a comprehensive diagnostic assessment, including digital impressions, photographs, and radiographs of the patient's dentition. These records are used to create a 3D digital model of the teeth, which serves as the foundation for the treatment plan.

For canine retraction, the orthodontist will design a series of aligners that apply specific forces to move the canines into the desired position. The treatment plan outlines the sequential movement of the teeth, with each aligner in the series designed to achieve a small, incremental change. Patients are instructed to wear each aligner for 1–2 weeks before progressing to the next set. Regular follow-up visits are scheduled to monitor progress and make any necessary adjustments to the treatment plan.

In cases where additional anchorage is needed, adjunctive treatments such as TADs or elastics may be used in combination with clear aligners. These auxiliary devices can enhance the effectiveness of aligner therapy, particularly for complex movements.

Self-ligating brackets

Self-ligating brackets are an innovative development in orthodontics, designed to improve the efficiency and comfort of tooth movement. Unlike traditional brackets, which use elastics or metal ties to hold the archwire in place, self-ligating brackets have built-in mechanisms (such as clips or doors) to secure the archwire. This system reduces friction and allows for smoother and more efficient tooth movement. This section discusses the mechanisms, advantages, limitations, and clinical applications of self-ligating brackets, with a focus on their role in canine retraction.^[6,25,26]

Mechanism

Self-ligating brackets operate using a passive or active mechanism to hold the archwire. Passive self-ligating brackets have a sliding mechanism that allows the archwire to move freely within the bracket, reducing friction. Active self-ligating brackets use a clip that applies pressure to the archwire, providing more control over tooth movement.

The reduction in friction with self-ligating brackets is a significant advantage. Traditional elastics or metal ties can create resistance between the archwire and bracket, slowing down tooth movement and requiring more force. In contrast, self-ligating brackets allow for smoother, more efficient movement by minimizing this resistance. This efficiency is particularly beneficial for canine retraction, where precise and controlled movement is necessary.

Clinical application

Self-ligating brackets are integrated into the orthodontic treatment plan based on the specific needs of the patient. For canine retraction, these brackets can be particularly advantageous due to their ability to reduce friction and apply more consistent forces. The orthodontist attaches the self-ligating brackets to the teeth and threads the archwire through the brackets. The built-in clips or doors secure the archwire, allowing it to move freely within the bracket slot.

This setup allows for efficient retraction of canines into the extraction spaces. The reduced friction and consistent force application help achieve controlled and precise tooth

movements. Additionally, the smaller profile of self-ligating brackets enhances patient comfort and reduces soft tissue irritation.

Self-ligating brackets can be used alone or in combination with other orthodontic appliances, such as TADs, to enhance anchorage and control. The orthodontist monitors the treatment progress through regular follow-up visits, making necessary adjustments to ensure optimal outcomes.

Piezoelectric surgery

Piezoelectric surgery is an innovative and minimally invasive technique used in orthodontics to enhance the efficiency of tooth movement, including the retraction of canines. This surgical method leverages ultrasonic vibrations to create precise cuts in the bone, reducing resistance to tooth movement and accelerating the remodeling process. Piezoelectric surgery is particularly beneficial for complex orthodontic cases where traditional methods may fall short in achieving desired outcomes.^[6,12,18]

Mechanism

Piezoelectric surgery utilizes ultrasonic vibrations generated by a piezoelectric device to make microincisions in the bone. These vibrations are highly controlled and focused, allowing for precise cutting without damaging the surrounding soft tissues, nerves, or blood vessels. The ultrasonic waves cause minimal trauma to the bone, which facilitates faster healing and reduces postoperative discomfort.

In the context of canine retraction, piezoelectric surgery can be used to create osteotomies (bone cuts) around the canines and the surrounding bone. These microincisions reduce the density of the bone and enhance its remodeling capacity, allowing for faster and more efficient tooth movement. The reduced bone resistance facilitates the retraction of canines into the desired position with greater control and predictability.

Clinical application

Piezoelectric surgery is typically performed in a clinical setting under local anesthesia. The orthodontist or oral surgeon uses a piezoelectric device to create precise microincisions in the bone around the canines and adjacent teeth. These incisions reduce bone resistance and facilitate the retraction of the canines.

For canine retraction, the procedure involves creating vertical and horizontal cuts in the alveolar bone to weaken its structure and promote faster remodeling. The reduced bone density allows the canines to move more easily into the extraction spaces. Following the surgery, orthodontic forces are applied using braces, elastics, or other appliances to guide the canines into their desired positions.

The postoperative care involves monitoring the healing process and ensuring that the patient follows proper oral hygiene practices. Regular follow-up visits are scheduled to assess progress and make any necessary adjustments to the orthodontic appliances.

COMPARATIVE ANALYSIS

Efficacy and efficiency

Traditional methods like elastic chains and NiTi coil springs are effective but may require frequent adjustments. Contemporary methods, such as TADs and self-ligating brackets, offer more controlled and efficient tooth movement. Clear aligners, while effective, depend heavily on patient compliance.^[1,8,11,25]

Patient comfort and compliance

Clear aligners and self-ligating brackets generally provide greater comfort and esthetic appeal, improving patient compliance. TADs, while effective, can cause discomfort due to their invasive nature. Traditional methods may be less comfortable due to the constant pressure applied by elastics and springs.^[21-25]

Clinical considerations

The choice of method depends on various factors, including the severity of malocclusion, patient preferences, and clinical expertise. A combination of methods may be employed to achieve optimal results. For instance, TADs can be used in conjunction with clear aligners or self-ligating brackets to enhance anchorage and control.^[6,15,20]

CONCLUSION

The evolution of canine retraction methods in orthodontics reflects ongoing advancements in materials and techniques. While traditional methods remain valuable, contemporary approaches offer enhanced control, efficiency, and patient comfort. Orthodontists must consider individual patient needs and clinical circumstances to select the most appropriate retraction method. Future research and technological developments will continue to shape the landscape of orthodontic treatment, providing even more effective solutions for canine retraction.

REFERENCES

- Proffit WR, Fields HW, Sarver DM. Contemporary Orthodontics. 5th ed. St. Louis: Mosby; 2012.
- Nanda R, Kuhlberg A. Biomechanics and Esthetic Strategies in Clinical Orthodontics. Netherlands: Elsevier Health Sciences; 2005.
- Papageorgiou SN, Zogakis IP, Papadopoulos MA. Clinical effectiveness of class II subdivision treatment: A systematic review and meta-analysis. *Eur J Orthod* 2014;36:547-59.
- Park HS, Kwon OW, Sung JH. Nonextraction treatment of an open bite with microcrew implant anchorage. *Am J Orthod Dentofacial Orthop* 2006;130:391-402.
- Keim RG, Gottlieb EL, Nelson AH, Vogels DS 3rd. 2008 JCO study of orthodontic diagnosis and treatment procedures. Part 1. Results and trends. *J Clin Orthod* 2008;42:625-40.
- Kulshrestha RS, Tandon R, Chandra P. Canine retraction: A systematic review of different methods used. *J Orthod Sci* 2015;4:1-8.
- Burstone CJ, Koenig HA. Optimizing anterior and canine retraction. *Am J Orthod* 1976;70:1-19.
- Ziegler P, Ingervall B. A clinical study of maxillary canine retraction with a retraction spring and with sliding mechanics. *Am J Orthod Dentofacial Orthop* 1989;95:99-106.
- Gjessing P. Biomechanical design and clinical evaluation of a new canine-retraction spring. *Am J Orthod* 1985;87:353-62.
- Lotzof LP, Fine HA, Cisneros GJ. Canine retraction: A comparison of two preadjusted bracket systems. *Am J Orthod Dentofacial Orthop* 1996;110:191-6.
- Dinçer M, Işcan HN. The effects of different sectional arches in canine retraction. *Eur J Orthod* 1994;16:317-23.
- Lewis PD. Canine retraction. *Am J Orthod* 1970;57:543-60.
- Shpack N, Davidovitch M, Sarne O, Panayi N, Vardimon AD. Duration and anchorage management of canine retraction with bodily versus tipping mechanics. *Angle Orthod* 2008;78:95-100.
- Kojima Y, Fukui H. Numerical simulation of canine retraction by sliding mechanics. *Am J Orthod Dentofacial Orthop* 2005;127:542-51.
- İşeri H, Kışınçisi R, Bzizi N, Tüz H. Rapid canine retraction and orthodontic treatment with dentoalveolar distraction osteogenesis. *Am J Orthod Dentofacial Orthop* 2005;127:533-41.
- Cernei ER, Chehab A, Olteanu DN, Romanec C, Panaite T, Zetu IN. Passive Self-ligating bracket systems: A scoping review of their claims regarding efficiency and effectiveness in orthodontic treatment. *Appl Sci* 2023;13:6322.
- Häsler R, Schmid G, Ingervall B, Gebauer U. A clinical comparison of the rate of maxillary canine retraction into healed and recent extraction sites--a pilot study. *Eur J Orthod* 1997;19:711-9.
- Wazwaz F, Seehra J, Carpenter GH, Papageorgiou SN, Cobourne MT. Duration of canine retraction with fixed appliances: A systematic review and meta-analysis. *Am J Orthod Dentofac Orthop* 2023;163:154-72.
- Nahidh M, Yassir YA, McIntyre GT. Different methods of canine retraction-Part 2. *J Baghdad Coll Dent* 2023;35:58-75.
- Idrees W, Kanwal L, Maaz M, Fida M, Sukhia RH. Comparison of canine retraction rate between miniscrew assisted micro-osteoperforation and conventional technique-a systematic review and meta-analysis. *Clin Invest Orthod* 2024;83:1-6.
- Sharma R, Soni DM. Temporary anchorage devices (TADs) in orthodontics: A review. *Asian J Res Med Med Sci* 2023;5:1-11.
- Baxi S, Bhatia V, Tripathi A, Dubey MP, Kumar P, Mapare S. Temporary anchorage devices. *Cureus* 2023;15:e44514.
- AlMogbel A. Clear aligner therapy: Up to date review article. *J Orthod Sci* 2023;12:37.
- Bichu YM, Alwafi A, Liu X, Andrews J, Ludwig B, Bichu AY, *et al.* Advances in orthodontic clear aligner materials. *Bioact Mater* 2023;22:384-403.
- Coronel-Zubiarte FT, Luján-Valencia SA, Meza-Málaga JM, Aguirre-Ipenza R, Echevarria-Goche A, Luján-Urviola E, *et al.* Effect of conventional and self-ligating brackets on periodontal health. Systematic review and meta-analysis. *J Clin Exp Dent* 2024;16:e358.
- Hendiani I, Prasetyo BC, Evangelina IA, Rizqita PA. The effects of using conventional and self-ligating brackets on oral hygiene and periodontal health status: A rapid review. *J Int Dent Med Res* 2023;16:384-93.