Review Article

Early Orthodontic Treatment: Past to Present

Yusaf C. Muhamedali*, Vaibhav Vashishta, Stuti Raj, Rajkumar Jaiswal, Shraddha Bind, Mrunali Kore
Department of Orthodontics, Sardar Patel Post Graduate Institute of Dental and Medical Sciences, Lucknow, Uttar Pradesh, India.

Abstract

The development of dental occlusion is a highly intricate process that spans from early intrauterine life to adulthood. The importance of early orthodontic intervention has been widely recognized as it aids in preventing severe dental malocclusions, reduces the need for complex treatments later in life, and enhances overall oral health and facial esthetics. Early orthodontic treatment (EOT) focuses on guiding the growth of jaw structures, correcting improper oral habits, and ensuring the proper alignment of primary and permanent teeth. The purpose of this comprehensive article is to explore the evolution of EOT from past to present, discussing the history, advancements, methodologies, benefits, challenges, and future directions in the field. Special emphasis will be placed on space management, habit correction, growth modification techniques, and digital orthodontics, among other aspects.

Keywords: Digital orthodontics, Early orthodontic treatment, Growth modification, Habit correction.

Introduction and Objectives

The field of orthodontics has evolved significantly over the years, with early intervention emerging as a key component in managing dental and skeletal malocclusions. Early orthodontic treatment (EOT) refers to the proactive measures taken during the primary or mixed dentition phases to prevent or mitigate the severity of developing malocclusions. This approach allows clinicians to take advantage of a child's natural growth potential, facilitating more effective and stable corrections compared to later interventions.

Malocclusions, if left untreated, can contribute to various complications, including impaired chewing and speech, increased susceptibility to dental caries and periodontal diseases, and adverse effects on facial esthetics and self-esteem.[1] Addressing these issues at an early stage can significantly reduce the complexity and duration of future orthodontic treatment, improving both functional and psychological outcomes for young patients.[2]

Historical Context

Orthodontics has a long and evolving history dating back to ancient civilizations. The early Egyptians, Greeks, and Romans documented rudimentary orthodontic techniques.[3] In the 18th century, Pierre Fauchard introduced dental appliances to correct malocclusion.[4] By the late 19th and early 20th centuries, Edward Angle revolutionized the field by developing a classification system for malocclusions and advocating for non-extraction treatment approaches.[5]

During the mid-20th century, the importance of early intervention became more apparent as orthodontists recognized that addressing skeletal and dental discrepancies at a young age could minimize the need for corrective surgeries later in life.[6]

With the advent of new materials, diagnostic tools, and biomechanical principles, EOT has become a standard practice in contemporary dentistry.[7]

The Science of Occlusal Development

Understanding occlusal development is fundamental in EOT. The occlusal development process consists of several stages:

1. Primary dentition stage (6 months–6 years): The initial eruption of primary teeth is crucial for speech development and mastication.[8]
2. Mixed dentition stage (6–12 years): The transition period where both primary and permanent teeth coexist. This is the most favorable stage for early intervention.[9]

Address for correspondence: Dr. Yusaf C. Muhamedali, Department of Orthodontics, Sardar Patel Post Graduate Institute of Dental and Medical Sciences, Lucknow, Uttar Pradesh, India.
Email: yousafali474@yahoo.com

Received: Mar. 24, 2025; **Accepted:** Apr. 20, 2025; **Published:** Apr. 27, 2025

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

How to cite this article: Muhamedali YC, Vashishta V, Raj S, Jaiswal R, Bind S, Kore M. Early Orthodontic Treatment: Past to Present. J Res Adv Dent 2025;16(3):1-3.

Access this article online

Website: www.jrad.co.in

DOI: https://doi.org/10.53064/jrad.2024.16.3.665

3. Permanent dentition stage (12+ years): The final stage where all permanent teeth have erupted and occlusal relationships become more defined.[10]

Failure to intervene during critical growth phases can result in malocclusions that require more extensive orthodontic procedures in later years.[11]

Space Management in Transitional Dentition

One of the primary goals of early orthodontic intervention is space management. Space maintenance and regaining are crucial for ensuring the proper eruption and alignment of permanent teeth.

Techniques for Space Maintenance

- Band and loop space maintainers: Prevent mesial migration of adjacent teeth after premature tooth loss[12]
- Distal shoe space maintainers: Used when the first molar has not yet erupted
- Lingual holding arch: Aids in preserving arch integrity in the mandibular arch
- Nance appliance: Prevents mesial drift of molars in the maxilla.

Techniques for Space Regaining

- Lip bumper appliance: Encourages distal movement of molars
- Headgear therapy: Assists in space regaining through extraoral forces
- Pendulum appliance: Aids in molar distalization for regaining lost space.

EOT Modalities

Preventive orthodontics

Preventive orthodontic treatment focuses on minimizing the risk of developing malocclusions. It includes:

- Space maintainers
- Habit-breaking appliances
- Serial extractions for guiding eruption.

Interceptive orthodontics

Interceptive orthodontic treatment is aimed at addressing early-developing malocclusions before they worsen.[13] Common interventions include:

- Expansion appliances for transverse deficiencies
- Functional appliances for Class II and III malocclusions
- Correction of anterior and posterior crossbites
- Early alignment of severely displaced teeth.

Corrective orthodontics

For cases where preventive and interceptive measures are insufficient, corrective orthodontic treatment, such as comprehensive fixed appliance therapy, may be required.

Growth Modification Techniques

Growth modification is an essential component of EOT, particularly in cases of skeletal discrepancies. By harnessing the child's natural growth potential, orthodontists can correct jaw alignment and facial proportions with minimal invasiveness.[14]

Common growth modification appliances:

- Twin block appliance: Encourages mandibular advancement for Class II malocclusion
- Reverse pull headgear: Used in Class III patients to stimulate maxillary growth
- Palatal expanders: Correct transverse discrepancies in cases of posterior crossbite
- Herbst appliance: Enhances mandibular positioning in growing patients.[15]

The Future of EOT

The future of early orthodontics is being shaped by advancements in biotechnology, digital technology, and patient-centered treatment approaches.

1. 3D-printed and custom appliances
   - Patient-specific brackets and aligners will enhance efficiency and reduce treatment time
   - Biocompatible materials will allow for improved durability and comfort.
2. Artificial intelligence in treatment planning
   - AI-driven diagnostics will enhance cephalometric analysis and predictive growth modulation
   - Real-time treatment adjustments based on patient progress will optimize results.
3. Teleorthodontics and remote monitoring
   - AI-assisted virtual consultations will improve accessibility and convenience
   - Wearable sensors in appliances will track treatment adherence.
4. Gene therapy and regenerative orthodontics
   - Genetic modulation of tooth eruption may allow for earlier and more effective interventions
   - Stem cell therapy could aid in craniofacial growth modification.
5. Minimally invasive techniques
   - Micro-osteoperforation and laser therapy will accelerate tooth movement
   - Piezoelectric bone stimulation may reduce treatment duration.

Discussion

The effectiveness of EOT has been widely studied, with most research indicating positive long-term outcomes. However, some controversies exist regarding the necessity and timing of intervention. While proponents argue that early treatment minimizes the complexity of later interventions, some studies suggest that certain cases may self-correct over time.

In addition, factors such as patient compliance, cost, and accessibility remain challenges in the widespread implementation of early orthodontic care.

Future research should focus on refining patient selection criteria to ensure that early interventions are recommended only when truly beneficial. Further exploration into genetic markers for predicting malocclusion severity and responsiveness to treatment could also enhance decision-making in orthodontics.

Conclusion

EOT offers significant advantages in terms of occlusal development, functional improvement, and esthetic enhancement. With the integration of digital technology and growth modification techniques, early intervention continues to evolve, providing better treatment outcomes with reduced complexity. Awareness and timely intervention remain key in ensuring optimal dental health for young patients.

By exploring the past, present, and future of early orthodontics, this article aims to provide a comprehensive understanding of its significance in contemporary orthodontic practice.

As research continues to advance, early orthodontics will remain an integral component of preventive and corrective dentistry, ultimately benefiting countless individuals by ensuring optimal dental development and overall oral health.

References

1. Proffit WR, Fields H, Larson B, Sarver DM. Contemporary Orthodontics. 6th ed. South Asia Edition-E-Book. Philadelphia, PA: Elsevier Health Sciences; 2019.
2. Bishara SE. Textbook of Orthodontics. New Delhi: Jaypee Brothers, Medical Publishers; 2001.
3. Moyers RE. Handbook of Orthodontics. 4th ed. Maryland: Year Book Medical Publishers; 1988.
4. Graber TM, Vanarsdall RL, Vig KW. Orthodontics: Current Principles and Techniques. 5th ed. United States: Mosby; 2012.
5. Little RM. The long-term stability of dental arch form. Am J Orthod Dentofac Orthop 1990;97:323-33.
6. Tulloch JC, Phillips C, Proffit WR. Benefit of early Class II treatment: Progress report of a two-phase randomized clinical trial. Am J Orthod Dentofac Orthop 1998;113:62-74.
7. Faltin K Jr., Faltin RM, Baccetti T, Franchi L, Ghiozzi B, McNamara JA Jr. Long-term effectiveness and treatment timing for Bionator therapy. Angle Orthod 2003;73:221-30.
8. Cobourne MT, editor. Orthodontic Management of Class II Malocclusion: An Evidence-Based Guide. Germany: Springer Nature; 2024.
9. Al Naqbi IA, Kaklamanos EG, Papadopoulou AK, Athanasiou AE. Orthodontic procedures, with or without extracting primary canines, for the interceptive management of palatally displaced permanent canines: a systematic review. J Dent Child (Chic) 2020;87(2):60-8.
10. King GJ, Keeling SD, Hocevar RA, Wheeler TT. The timing of treatment for Class II malocclusions in children: A literature review. Angle Orthod 1990;60:87-97.
11. Proffit WR, Tulloch JC. Preadolescent Class II problems: Treat now or wait? Am J Orthod Dentofac Orthop 2002;121:560-2.
12. Graber TM, Rakosi T, Petrovic AG. Dentofacial Orthopedics with Functional Appliances. United States: Mosby; 1985.
13. Carlson DS. Biological rationale for early treatment of dentofacial deformities. Am J Orthod Dentofacial Orthop 2002;121:554-8.
14. Gomes AS, Lima EM. Mandibular growth during adolescence. Angle Orthod 2006;76:786-90.
15. Pancherz H. The Herbst appliance--its biologic effects and clinical use. Am J Orthod 1985;87:1-20.