

Comparison of Curcumin Gel with Triamcinolone Acetonide in Treatment of Recurrent Aphthous Stomatitis

Anjani Kumar Shukla^{1*} Ashok Kumar² Ravi Kumar³ Rakhi Sinha⁴

¹Tutor, Department of Oral Medicine and Radiology, Dental Institute, Rajendra Institute of Medical Sciences, Ranchi, India.

²Senior Lecturer, Department of Pedodontics and Preventive Dentistry, Vananchal Dental College and Hospital, Garhwa, Jharkhand, India.

³Assistant Professor, Department of Dentistry, Heritage Institute of Medical Sciences, Varanasi, Uttar Pradesh, India.

⁴Senior Lecturer, Department of Oral Medicine and Radiology, Vananchal Dental College and Hospital, Garhwa, Jharkhand, India.

ABSTRACT

Aim: To assess and compare the efficacy of Curcumin with Triamcinolone acetonide in the gel form in treatment of minor RAS.

Materials and Methods: As a randomized clinical trial, 60 patients of either sex with clinically diagnosed RAS were randomly divided into 2 groups Curcumin gel group (Group I) and Triamcinolone Acetonide gel group (Group II). Patients in each group were asked to apply the gel three times a day on each ulcer. estimation of efficacy of gel was done on the basis of time required for regression in pain, size, and number of the ulcers.

Results: The results showed significant difference in size, pain, number, and duration of ulcers in Group I and Group II within a period of 7 days. However, no significant difference was noted in both the groups in the treatment of RAS.

Conclusion: Curcumin has strong antioxidant, with anti-inflammatory, immunomodulatory and analgesic properties. It can be used as an effective alternative to steroids in treatment of RAS.

Keywords: Curcumin, recurrent aphthous stomatitis, triamcinolone acetonide.

INTRODUCTION

Recurrent aphthous stomatitis (RAS), commonly known as aphthous ulcer, is an most common disease occurring in oral cavity. It is characterized by the development of painful, recurring solitary, or multiple ulcers in the oral cavity. The ulcers are frequently multiple, small, round, or ovoid, with well circumscribed margins, having yellow or grayish floor and are surrounded by erythematous base.[1]

RAS is classified into 3 types:

1. Minor RAS: Ulcers are less than 1 cm in size. Ulcers heal within 10-14 days without scarring.

2. Major RAS also known as Sutton's disease. Ulcers more than 1 cm in diameter. The ulcers persist for up to 6 weeks and heal with scarring.

3. Herpetiform ulceration is characterized by recurrent multiple ulcers, may be up to 100 in number. These are small in size. Lesions may coalesce to form large ulcers. These ulcers lasts for about 10-14 days.[1]

RAS is a multifactorial disease which occur due to bacterial infections like infection with *Streptococcus sanguis*, immunologic abnormalities, nutritional deficiencies like iron, vitamin B12, folic acid deficiency, and certain precipitating factors like trauma, endocrine conditions, psychic, and allergic factors. The lesion is self-limited, painful, high frequency of occurrence, and multifactorial etiology.

Received: Feb. 19, 2019; Accepted: Apr. 25, 2019

*Correspondence Dr. Anjani Kumar Shukla.

Department of Oral Medicine and Radiology, Dental Institute, Rajendra Institute of Medical Sciences, Ranchi, India.

Email: anjanisukla@yahoo.com

So an well-organized therapeutic strategy is needed to provide relief to the patients.[2-4]

In Current treatment modalities include use of topical steroids like Triamcinolone acetonide gel, which is a fluoride synthetic corticosteroid. As corticosteroids are known to get absorbed systemically even on topical application. So these days herbal medicines are gaining more interest due to their decreased side effects.[4-6]

Curcumin is a principle spice in India. It is bright yellow in color and is known from centuries for its strong antioxidant, antiseptic, antibacterial, anti-inflammatory, immunomodulatory, and analgesic properties. The anti-inflammatory properties of curcumin may be certified to its ability to inhibit both biosynthesis of inflammatory prostaglandin which further blocks cyclooxygenase and lipooxygenase activity, thereby inhibiting prostaglandin release and neutrophil function during inflammatory stage.[7-10]

So considering the above background, the aim & objectives of this study was to assess and compare the efficacy of Curcumin with Triamcinolone acetonide in the gel form in treatment of RAS.

MATERIALS AND METHODS

The study was a randomized, double blind, clinical trial and was carried out over a period of 6 months, from may 2017 to November 2017, After obtaining the written informed consent, 60 patients of either sex with clinically diagnosed minor RAS were selected from among the out patients visiting the Department of Oral Medicine and Radiology. Patients willing to participate in the study and with clinically diagnosed minor RAS noticed were included in the study.

Patients were excluded from the study if the ulcers were present at inaccessible areas, if they had major RAS, traumatic ulcer, denture stomatitis or if they were suffering from any systemic illness. The trial was approved by the research and ethical committee of vananchal dental college and hospital, Garhwa, Jharkhand.

Method of collection of data

Detailed case history, extraoral, and intraoral examination was performed on the patients, and 60

patients with clinically diagnosed minor RAS were selected for the study based on inclusion and exclusion criteria. These 60 patients were randomly divided into two group:

Group I 30 patients (Curcumin gel)

Group II 30 patients (Triamcinolone Acetonide gel)

Ulcer size was measured along its longest diameter using a sterile vernier caliper and the pain score was noted from 0-10, using the Visual Analogue Scale (VAS). Patients were asked to apply the gel three times a day on each ulcer after meals and do not take eat food or water for half an hour after application of the gel. All patients were provided with same measuring applicator and were instructed about the quantity and method of gel application on the first appointment itself. Patients were examined on day 0, 3, 5, and 7 to check for regression number and/or size of ulcer and any discomfort due to the gel. Pain score was noted daily by the patient.

Clinical estimation of ulcer

The response of treatment was assessed by quantitative measurement of diameter of ulcer using vernier caliper, at each follow-up visits by a single observer. The patients were instructed to note the pain score every morning on his/her record card as per the VAS. Assessment of efficacy of gel was done on the basis of time required for regression in pain, size and number of the ulcer.

Details of Curcumin gel:

- Curenex oral gel (Abbott Pharmaceuticals).
- Per gram of gel contained 10 mg of curcuma longa extract.

Details of Triamcinolone Acetonide gel:

- Kenacort oral paste (Abbott Pharmaceuticals).
- The gel contains 0.1% triamcinolone acetonide.

Data was presented as mean $\pm$ standard deviation (SD) for quantitative variables with normal distribution and significance of difference was accepted at $P < 0.05$. Background and demographic data were summarized with descriptive statistics. To evaluate the efficacy, Mann-Whitney U test was used and statistical analysis was done using Statistical Package for the Social Sciences (SPSS) 18 software.

RESULTS

Total of 60 patients [Table 1] participated in the study and all patients completed the study protocol.. No significant difference was identified between patients randomly assigned to the group I or group II with regard to basic demographic data including age and gender; size and of ulcers; and pain score on day 0. The reduction in pain was statistically significant in both the groups from day 0-day 7, respectively; but on comparison between group I and group II, no statistical significance was noted in the pain score. [Table 2]

The reduction in size and number of ulcers were statistically significant in both the groups from day 0-day 7 (noted on day 0, 3, 5, 7), respectively; but on comparison between group I and group II, no statistical significance was noted in the size and number of ulcers. There was statistical significance in duration of ulcer in both the groups from day 0-day 7, respectively; but on comparison between group I and group II, no statistical significance was noted in the duration of ulcers. [Table 3]

DISCUSSION

The multifactorial etiology of RAS and lack of effective treatment modality makes it a challenge for the dentist to relieve the patient from the morbidity associated with RAS. As the treatment of RAS aims at providing symptomatic relief for the patient, the goals of treatment include reduction in healing time of ulcer, decrease pain, size, and number of ulcers.[4-6] Current treatment modalities include use of systemic and topical steroids, cauterization, antibiotics, mouth rinses containing active enzymes, laser treatments, and combination therapy.[9]

As curcumin is known for its strong antioxidant, antiseptic, antibacterial, anti-inflammatory, immunomodulatory, and analgesic properties, a study was carried out to evaluate the therapeutic effect of viscous curcumin solution at 10% and 50%, on the healing of ulcer in patients with RAS showed a good percentage of complete healing of ulcer in patients with RAS.[11,12]

In 2012, an randomized placebo controlled trial was done using 2% curcumin gel in treatment of minor RAS, which showed that it is a well-tolerated

effective antibacterial, anti-tumor agent with potent anti-inflammatory and analgesic properties.[13]

This study showed that there was significant difference between the pain score, size, number, and duration of ulcers. Statistical significance was not noted between all the parameters when both the groups were compared with each other. The results of this study, as consistent with other studies provide positive evidence that curcumin gel can be used as an effective and safer alternative to steroids in treatment of minor RAS.

CONCLUSION

The topical application of curcumin and triamcinolone acetonide gel show similar efficiency in the treatment of minor RAS. The results obtained from the present study provide strong evidence that curcumin gel can be used as an useful and safer alternative to steroids in treatment of RAS.

CONFLICTS OF INTEREST

The authors declare they have no potential conflict of interests regarding this article.

REFERENCES

1. Preeti L, Magesh K, Rajkumar K, Karthik R. Recurrent aphthous stomatitis. J Oral Maxillofac Pathol 2011;15:252-6.
2. Greenberg MS, Glick M, Ship JA. Burket's Oral Medicine. Hamilton: BC Decker Inc; 2008. Volume 11. p. 57-60.
3. Porter SR, Hegarty A, Kaliakatsou F, Hodgson TA, Scully C. Recurrent aphthous stomatitis. Clin Dermatol 2000;18:569-78.
4. Ship JA. Recurrent aphthous stomatitis. Oral Surg Oral Med Oral Pathol Oral Radiol Endod 1996;81:141-7.
5. Akintoye SO, Greenberg MS. Recurrent aphthous stomatitis. Dent Clin North Am 2005;49:31-47.
6. Shashy RG, Ridley MB. Aphthous ulcers: A difficult clinical entity. Am J Otolaryngol 2000;21:389-93.
7. Akram M, Shahab-Uddin, Ahmed A, Khan U, Hannan A, Mohiuddin E, et al. Curcuma longa and

curcumin: A review article. Rom J Biol Plant Biol 2010;55:65-70.

8. Chainani-Wu N. Safety and anti-inflammatory activity of curcumin: A component of turmeric (*Curcuma Longa*). J Altern Complement Med 2003;9:161-8.

9. Thorne R. *Curcuma longa* (Turmeric). Altern Med Rev 2001;6:62-8.

10. Janson MD. Curcumin. Healthy Living 2000;2:1-3.

11. Itsuki K, Tetsuya F, Teruo N, Toshihiko I, Keinosuke N. Therapeutic effect of curcumin

chewable on aphthous ulcer and traumatic ulcer. Clin All-Round 2006;55:204-8.

12. Al-Saffar MT. The therapeutic effect of viscous solution of curcumin in the treatment of recurrent aphthous stomatitis (RAS). Al-Rafidain Dent J 2006;6:48-52.

13. Manifar S, Obwaller A, Gharehgozloo A, Boorboor Shirazi Kordi HR, Akhondzadeh S. Curcumin gel in treatment of minor aphthous ulcer: A randomized, placebo-controlled trial. J Med Plants 2012;11:41.