

Visual Treatment Objective: A Comprehensive Overview

Saif Alam*, Jitendra Bhagchandani, Vaibhav Vashishta, Amit Kumar Singh, Tushant Rastogi, Ayushi Singh

Department of Orthodontics, Sardar Patel Post Graduate Institute of Dental and Medical Sciences, Lucknow, Uttar Pradesh, India.

Abstract

The visual treatment objective prediction tracing is a crucial instrument in the diagnosis and treatment of dentofacial and craniofacial abnormalities. It facilitates the evaluation of facial esthetics and the planning of necessary surgical interventions and adjustments. The visualization of the treatment objective can be a straightforward but reasonably accurate way to forecast the relationship between the incisor and molar. It may also be used to assess whether any “midcourse” adjustments are made during treatment and to compare the outcome to the initial forecast. The visual representation of the patient’s progress is crucial in guiding the therapist to evaluate and track therapy advancement and implement strategic modifications. A significant determinant in forecasting the appropriate treatment for a patient is the clinician’s personal assessment and appraisal, encompassing both the physical condition and mental disposition.

Keywords: Cephalometric analysis, Cephalometric tracings, Forecast, Growth prediction.

INTRODUCTION AND OBJECTIVES

The visual treatment objective (VTO) is an indispensable diagnostic tool in orthodontics that helps predict treatment outcomes by analyzing the dentoskeletal framework.^[1] It allows orthodontists to forecast how growth and treatment will affect incisor and molar positions, as well as the relationship between skeletal structures and soft tissues.

VTO establishes achievable goals from the outset and provides continuous guidance for treatment progression and final results. By incorporating cephalometric tracings, orthodontists can visualize tooth movement, skeletal changes, and their impact on facial esthetics.^[2]

Key objectives include

- Forecasting growth and changes over the treatment period
- Evaluating soft-tissue structures and facial balance
- Assessing arch length discrepancies and determining treatment mechanics
- Identifying candidates for surgical and surgical-orthodontic procedures
- Comparing treatment outcomes to initial predictions to guide midcourse corrections.

HISTORICAL CONTEXT

The VTO concept was pioneered in 1948 by Dr. William B. Downs, who developed methods for predicting facial

changes during orthodontic treatment.^[3] He emphasized cephalometric analysis, particularly the role of the cranial base and mandibular plane. Robert Ricketts expanded on Downs’ work by integrating cranial base references and mandibular growth analysis.^[4]

Key contributions include

- Ricketts’ analysis: Introduced facial axis and maxillary growth prediction
- Holdaway’s method: Integrated soft-tissue considerations, focusing on lip strain and harmony
- Katona’s approach: Applied mathematical precision to cephalometric analysis, optimizing relationships between skeletal and dental structures.

Over time, advancements in imaging and software have further refined VTO applications, making it an essential tool for modern orthodontics.^[5]

Advantages of VTO

The VTO provides several benefits, enhancing both diagnosis and treatment:

Address for correspondence: Dr. Saif Alam,
Department of Orthodontics, Sardar Patel Post Graduate Institute of Dental
and Medical Sciences, Lucknow, Uttar Pradesh, India.
Email: amu_saifalam91@live.co.uk

Received: Jan. 15, 2025; **Accepted:** Feb. 16, 2025 **Published:** ***

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

How to cite this article: Alam S, Bhagchandani J, Vashishta V, Singh AK, Rastogi T, Singh A. Visual Treatment Objective (VTO): A Comprehensive Overview. *J Res Adv Dent* 2025;16:(2):1-3.

Access this article online

Website:
www.jrad.co.in

DOI:
<https://doi.org/10.53064/jrad.2024.16.2.652>

1. Predictive capabilities: Accurately forecasts changes in dentoskeletal structures and soft-tissue profiles
2. Differentiates growth versus treatment effects: Distinguishes between changes due to natural growth and therapeutic interventions
3. Improves treatment planning: Identifies challenges, suggests alternative strategies, and helps plan anchorage and tooth movement
4. Ensures continuity and precision: Continuous evaluation through superimpositions ensures alignment with treatment goals
5. Enhances patient communication: Visual aids improve patient understanding and motivation
6. Educational resource: Serves as a valuable teaching tool for clinicians and students.

The ability to overlay VTO tracings on initial records and compare predicted versus achieved outcomes provides orthodontists with invaluable feedback for treatment adjustments and learning.^[6]

Types of VTO

The VTO framework encompasses several categories based on treatment goals and methodologies:

Cephalometric VTO^[1]

- a. Rickett's method: Focuses on cranial base, mandibular growth, and maxillary expansion predictions while incorporating soft-tissue analysis.^[7] (Figure 1 depicts the landmarks used in Rickett's analysis)
- b. Holdaway's method: Emphasizes soft-tissue balance, lip strain, and facial esthetics.^[8] (Figure 2 depicts the landmarks used in Holdaway's analysis)
- c. Katona's method: Uses geometric principles to optimize relationships between incisors and skeletal angles. (Figure 3 depicts Katona's method)




Figure 1: Landmarks used in Rickett's method

- d. Lionel Sadowsky method: Involves cephalometric analysis with films taken in a lips-closed position, even under slight pressure.^[9] Key reference lines include the Basion–Nasion (BaN) for cranial base relationships and Nasion to point A (NA) for maxillary position
- e. Mini-VTO: The mini-VTO consists of three key phases:
 - Creating a cephalometric record and documenting standard diagnostic measurements
 - Calculating a 2-year prediction for the position of the upper and lower jaws, considering growth and treatment effects
 - Aligning the teeth optimally within the predicted skeletal framework.

The eleven-factor summary analysis

The eleven-factor summary evaluates key cephalometric parameters, including facial axis, mandibular plane angle,




Figure 2: Landmarks used in Holdaway's method




Figure 3: Katona's method




Figure 4: Landmarks used in TOMAC method. TOMAC: Treatment outcome and management analysis by computer

and lower facial height. These measures help guide treatment planning, anchorage decisions, and extraction needs.^[10]

Surgical VTO

Surgical VTO evaluates treatment outcomes that combine orthodontics and surgical interventions. Techniques include:

- Cadei and Bassini method: Assesses facial asymmetry and skeletal changes^[10]
- Donald Gay method: Specializes in tracing asymmetric cases^[11]
- Treatment outcome and management analysis by computer (TOMAC) analysis: Focuses on vertical and horizontal dimensions of soft tissue and skeletal structures.^[12] (Figure 4 depicts the landmarks used in TOMAC analysis)
- Video cephalometric diagnosis: Visualizes complex craniofacial deformities to guide surgical planning.

Dental VTO

Dental VTO predicts tooth movement and alignment within the arch.^[13] (Figure 5 depicts the method of Dental VTO), including:

- Evaluation of arch length discrepancies
- Determining optimal positions for incisors and molars
- Adjusting appliance settings for precise outcomes.

Functional VTO

Functional VTO focuses on the relationship between growth patterns, orthopedic appliances, and facial structure corrections.^[14] It is particularly effective for:

- Correcting Class II or III malocclusions




Figure 5: Dental VTO

- Managing open bites and crossbites
- Influencing mandibular growth in young patients.

Evaluation

Evaluation in VTO relies on superimpositions of tracings to measure changes in skeletal, dental, and soft tissue structures.^[14]

The process involves five superimposition areas:

- Chin and mandibular plane: Tracks mandibular growth and chin positioning
- Maxilla: Assesses maxillary development and point A changes
- Occlusal plane: Evaluates bite alignment and occlusal adjustments
- Teeth in mandible and maxilla: Tracks incisor and molar positioning
- Soft-tissue profile: Monitors esthetic outcomes and facial balance.

DISCUSSION

The integration of VTO into treatment planning improves:

- Predictability of outcomes by combining growth predictions with mechanical interventions
- Patient understanding and acceptance of treatment goals through visual simulations
- Treatment efficiency by highlighting potential complications and alternative strategies early in the planning process.

Despite its strengths, VTO relies heavily on the clinician's expertise and accurate data input.^[15] Unpredictable growth patterns or treatment responses may impact its accuracy, requiring iterative adjustments during treatment.

CONCLUSION

VTO remains a powerful tool in orthodontics, combining cephalometric methods, surgical tracings, and functional analyses to ensure holistic treatment outcomes. By addressing skeletal, dental, and soft-tissue factors, VTO enables orthodontists to deliver precise, esthetic, and functional results. With advancements in imaging and diagnostic tools, its applications continue to evolve, reinforcing its role in modern orthodontic practice.

REFERENCES

1. Bench RW, Gugino CF, Hilgers JJ. Bio-progressive therapy. *J Clin Orthod* 1977;11:661-71, 674-82 contd.
2. Holdaway RA. A soft-tissue cephalometric analysis and its use in orthodontic treatment planning. Part I. *Am J Orthod* 1983;84:1-28.
3. Jacobsen A. A visualized treatment objective. *J Clin Orthod* 1980;14:554-71.
4. Magness WB. The mini-visualized treatment objective. *Am J Orthod Dentofacial Orthop* 1987;91:361-74.
5. Lopez FJ. Diagnosis with computer-aided drafting. *J Clin Orthod* 1986;20:327-9.
6. Guess MB, Solzer WV. Computer treatment estimates in orthodontics and orthognathic surgery. *J Clin Orthod* 1989;23:262-8.
7. Sarver DM. Video cephalometric diagnosis (VCD): A new concept in treatment planning? *Am J Orthod Dentofacial Orthop* 1996;110:128-36.
8. Brandt S, Ricketts RM. Ricketts on growth prediction. *J Clin Orthod* 1975;9:340-62, 352-62.
9. Ricketts RM. Bioprogressive therapy as an answer to orthodontic needs. Part I. *Am J Orthod* 1976;70:241-68.
10. Servoss JM. Derivation of acceptable arrangements in the Steiner analysis. *Angle Orthod* 1971;41:146-9.
11. Tweed CH. The Frankfort-mandibular plane angle in orthodontic diagnosis, classification, treatment planning, and prognosis. *Am J Orthod Oral Surg* 1946;32:175-230.
12. Moshiri F, Jung ST, Sclaroff A, Marsh JL, Gay WD. Surgical diagnosis and treatment planning: A visual approach. *J Clin Orthod* 1982;16:37-59.
13. McLaughlin RP, Bennett JC. The dental VTO: An analysis of orthodontic tooth movement. *J Clin Orthod* 1999;33:394-403.
14. Johnston LE. A statistical evaluation of cephalometric prediction. *Angle Orthod* 1968;38:284-304.
15. McCollum TG. TOMAC: An orthognathic treatment planning system. Part 1 soft-tissue analysis. *J Clin Orthod* 2001;35:356-64.