

Radix: A Clinical Challenge for Endodontists

Nutan Mala¹ Smita^{2*} C. K. Wang³¹PG Student, Department of Conservative Dentistry and Endodontics, Buddha Institute of Dental Sciences & Hospital, Patna, Bihar, India.²PG Student, Department of Conservative Dentistry and Endodontics, Buddha Institute of Dental Sciences & Hospital, Patna, Bihar, India.³PG Student, Department of Conservative Dentistry and Endodontics, Buddha Institute of Dental Sciences & Hospital, Patna, Bihar, India.

ABSTRACT

Background: Mandibular molars can have an additional root located lingually (the radix entomolaris) or buccally (the radix paramolaris). If present, awareness and understanding of this unusual root and its root canal morphology can contribute to the successful outcome of root canal treatment. This report discusses the endodontic treatment of a mandibular molar with a radix paramolaris.

Keywords: Radix paramolaris, unusual morphology, endodontic treatment.

INTRODUCTION

According to Swartz, Skidmore and Griffen, mandibular first molars have a significantly lower success rate compared with other teeth¹. Missed canals and the failure to remove all the microorganisms and pulp remnants from the root canal system are probably the main reasons for persistent infection around endodontically- treated molars². It is therefore important that clinicians have an awareness and good understanding of the variations in root canal morphology of the mandibular first molar. Permanent mandibular first molars in Caucasian populations are generally two rooted teeth (one mesial and one distal root) with two mesial and one distal root canals^{3 4}. The two mesial root canals can end up in two distinct apical foramina, or they can merge together at the root tip end into one apical foramen⁵. According to Calberson *et al.* (2007) the aetiology behind the formation is still unclear, but it could be related to external factors during odontogenesis. Racial genetic factors can also influence the profound expression of a particular gene that can result in the more pronounced phenotypic manifestation⁶.

Fig 1: Classification (de moor)

Types	Features
Type I	No curvature
Type II	Curvature in the coronal third and straight continuation to the apex
Type III	Curvature in the coronal third and additional buccal curvature from the middle third to the apical third of the root.
Small type	Root length less than half that of the distobuccal root.
Conical type	Cone-shaped extension with no root canal



Fig 2: Classification (Carlsen and Alexandersen).

Received: May. 16, 2020; Accepted: July. 29, 2020

*Correspondence Dr. Smita.

Department of Conservative Dentistry and Endodontics, Buddha Institute of Dental Sciences & Hospital, Patna, Bihar, India.

Email: smitapdch@gmail.com

Carlsen and Alexandersen's classification of RE. (a & b) Types A and B refer to a distally located cervical part of the RE with two normal and one normal distal root components, respectively, (c) Type C refers to a mesially located cervical part, (d) AC refers to a central location, between the distal and mesial root component.

CASE REPORT

A 45-year-old female patient was referred to the Department of Conservative Dentistry and Endodontics for endodontic treatment of the mandibular left first molar. On clinical examination, the tooth was decayed and showed signs of irreversible pulpitis. Radiographic examination showed the presence of a third root, which is indicative of Radix (Fig. 3)




Fig 3: IOPA

Further a CBCT scan was done to verify the position of the radix. The CBCT scan revealed that the root is present on the buccal side which is indicative of Radix Paramolaris.(Fig.4)




Fig 4: CBCT of the affected tooth.

After the location of the canal was assessed, the access preparation was done under rubber dam with an endo-access bur, and canal orifice was located with DG 16 endodontic explorer. Initial negotiation of the root canals was confirmed with K- file 15. The fourth canal (disto buccally) was located between mesial and distal root canal orifice, more nearer to distal orifices (Fig.5)




Fig 5: Showing the location of the fourth canal.

The canal lengths were determined radiographically with K file ISO 15 size and electronically with Dentaport Root ZX (J Morita). Cleaning was done with 17% EDTA and 3% sodium hypochlorite and shaped with Protaper Gold rotary system till the size of F-2. A temporary calcium hydroxide paste and temporary filling were put in place. The root canals were filled with gutta-percha and AH26 (De Trey Dentsply, Konstanz, Germany). The opening cavity was sealed with glass ionomer cement Fuji IX (GC Corp., Tokyo, Japan). This was confirmed with the help of an IOPA Radiograph (Fig.6)



Fig 6: Post-operative Radiograph.

Case 2

A male patient came to the Department of Conservative Dentistry and Endodontics, with a complaint of pain in the lower left back tooth region. On clinical examination, the tooth was

decayed and showed signs of irreversible pulpitis. Radiographic examination of which showed the presence of an extra root which is indicative of radix (Fig . 7). Access opening was done under rubber dam after anaesthetizing the tooth with an endo-access bur, and canal orifice was located with DG 16 endodontic explorer. Initial negotiation of the root canals was confirmed with K- file 15. The extra canal was found more nearer to the distal orifice. The canal lengths were determined radiographically with K file ISO 15 size and electronically with Dentaport Root ZX (J Morita) (Fig 8).

Cleaning was done with 17% EDTA and 3% sodium hypochlorite and shaped with Protaper Gold rotary system till the size of F-2. Master cone radiograph revealed the proper fitting of cones (Protaper GP-DENTSPLY). The cone fit, with radiographic exposure taken at 300 mesial angulation confirmed the presence of an RE.



Fig 7: Preoperative radiograph.




Fig 8: Postoperative radiograph.

Radix entomolaris, first described by Carabelli, is an anatomical variant in the first permanent mandibular molar typically characterized by an additional third root located distolingually. RE occurs in first, second and third molars with the lowest prevalence in second mandibular molars^{7 8 9}. Although the presence of RE differs within associated ethnical groups, it should be regarded as a normal anatomical variation within the Mongoloid population. Studies show no significant predilection for gender or side distribution with bilateral occurrence ranging between 50-67%¹⁰. Understanding the complexity of the anatomical variants seen in the first permanent mandibular molar proves to be invaluable in the clinical approach when treating these cases endodontically. Matherne *et al.* (2008) showed that CBCT images result in the identification of a greater amount of root canal systems in teeth compared to conventional radiography. The study also concluded that the combination of CBCT scanning with the dental operating microscope was an important diagnostic tool for locating and identifying root canals¹¹.

Protaper Gold (Dentsply/Maillefer) was used for root canal preparation for most of the cases depicted in this article. The key benefits of Protaper Gold include simplicity, excellent cutting efficiency and predictable final canal shape to allow for cone fit with tug-back. Another benefit of the system is the fact that the instrument is manufactured from M-wire and not traditional nickel-titanium alloy. Research by Johnson *et al.* (2008) demonstrated that the M-wire alloy could reduce cyclic fatigue by 400% compared with similar instruments manufactured from conventional nickel-titanium alloys.^{4,6} The added metallurgical benefit contributes towards more flexible instruments, increased safety and protection against instrument fracture¹².

CONCLUSION

The successful outcome of root canal treatment depends to a large extent on access, cleaning and shaping and three-dimensional obturation of the entire root canal system. CBCT technology, as well as proper angulation when acquiring radiographic images, proves helpful in locating canals in especially first mandibular molars with a high

incidence of anatomical variations. A thorough understanding of the prevalence of RE, its anatomical variations, as well as a radiographic and clinical diagnosis will provide the clinician with a better understanding of its complexity in order to ensure successful treatment outcomes.

CONFLICT OF INTEREST

No potential conflict of interest relevant to this article was reported.

REFERENCES

1. Swartz DB, Skidmore A, Griffin J. Twenty 1. years of endodontic success and failure. *J Endod.* 1983;9(5):198-202.
2. Himel V, McSpadden J, Goodis H, Cohen S, Hargreaves K. *Pathways of the pulp*, ninth edition. 2006.
3. Barker B, Parsons K, Mills P, Williams G. Anatomy of root canals. III. Permanent mandibular molars. *Aus Dent J.* 1974;19(6):408-13.
4. Vertucci FJ. Root canal anatomy of the human permanent teeth. *Oral surg, Oral Med, Oral Pathol.* 1984;58(5):589-99.
5. Vertucci F, Williams R. Root canal anatomy of the mandibular first molar. *JNJ Dental Assoc.* 1973;45(3):27-8 passim.
6. Rebeiro F, Consolaro A. Importancia clinica y antropologica de la raiz distolingual en los molars inferiores permanentes. *Endodoncia.* 1997;15:72-8.
7. Tu M-G, Huang H-L, Hsue S-S, Hsu J-T, Chen S-Y, Jou M-J, et al. Detection of permanent three-rooted mandibular first molars by cone-beam computed tomography imaging in Taiwanese individuals. *J Endod.* 2009;35(4):503-7.
8. Segura-Egea JJ, Jimenez-Pinzon A, Rios-Santos JV. Endodontic therapy in a 3-rooted mandibular first molar: importance of a thorough radiographic examination. *J Can Dent Assoc.* 2002;68(9):541-5.
9. Garg AK, Tewari RK, Kumar A, Hashmi SH, Agrawal N, Mishra SK. Prevalence of three-rooted mandibular permanent first molars among the Indian population. *J Endod.* 2010;36(8):1302-6.
10. Steelman R. Incidence of an accessory distal root on mandibular first permanent molars in Hispanic children. *ASDC J Dent Child.* 1985;53(2):122-3.
11. Matherne RP, Angelopoulos C, Kulild JC, Tira D. Use of conebeam computed tomography to identify root canal systems in vitro. *J Endod.* 2008;34(1):87-9.
12. Gutmann J, Gao Y. Alteration in the inherent metallic and surface properties of nickel-titanium root canal instruments to enhance performance, durability and safety: a focused review. *Int Endod J.* 2012;45(2):113-28.